

Likelihoods on coalescents: a Monte Carlo sampling approach to inferring parameters from population samples of molecular data¹

BY JOSEPH FELSENSTEIN, MARY K. KUHNER, JON YAMATO, AND PETER BEERLI

*Department of Genetics, University of Washington, Box, 7360, Seattle WA
98195-7360*

² When population samples of molecular data, such as sequences, are taken, the members of the sample are related by a gene tree whose shape is affected by the population processes, such as genetic drift, change of population size, and migration. Genetic parameters such as recombination also affect that genealogy. Likelihood inference of these parameters involves summing over all possible genealogies. There are a vast number of these, so that exact computation is not possible. Griffiths and Tavaré have proposed computing these likelihoods by Monte Carlo integration. Our group is doing this by the Hastings-Metropolis method of Markov Chain Monte Carlo integration. We now have, in our LAMARC package, programs to do this for constant-sized and growing populations, and for geographically structured populations. The bias of the estimator of population growth rate is discussed. One can also allow for samples stratified in time, as with fossil DNA or sequential samples from the population of virus in a patient. A program for recombining sequences is in progress, and we hope to put together an object-oriented environment which can cope with a variety of evolutionary forces.

1. Introduction. Samples of genes from natural populations of organisms are related by a genealogy, which is usually unknown. At the level of the copies of the genes, such a genealogy would specify where each copy of the gene came from. Thus, a particular copy that we sample may have come from the mother of that individual, from her father, from his father, from his mother, and so on, back in time. Other copies are doing the same. As we go back, occasionally two of these lineages will coalesce, as it happens that two copies of a gene are descended from the same parental copy. Thus, my great-great-great-grandmother might happen to be the sibling of your great-great-great-grandfather, and the genes we possess might then turn out to be copied from the same copy in one of their parents. Such coalescences are inevitable in natural populations.

Figure 1 shows such a pattern of ancestry. Each circle is an individual who has two copies of the gene; we are concerned not just with the genealogy of the individuals, but with the genealogy at the gene level. In the figure, time

¹Supported by NIH grant no. GM-51929 and NSF grant no. BIR-9527687

²AMS 1991 subject classifications. Primary ; secondary

Key words and phrases. Coalescent; Molecular evolution; genetics




FIG. 1. A coalescent tree of gene copies that is formed in a diagram showing from which gene in the previous generation each gene copy comes. Large circles are individuals, small circles are copies of genes. Three copies in the current generation trace back to two copies 6 generations earlier.

flows upwards. The sample consists of three copies of the gene taken from the latest generation (at the top). Arrows show the copies of the gene transmitted from parent to offspring. When we go backwards in time along the arrows, we go downwards, and the lineages gradually coalesce. The rate of this coalescence is higher in small populations than in large ones, simply because the chance that the ancestors of two copies of the gene are the same is greater in a small population.

We have reasonably straightforward models of change in the DNA sequences of such genes, based on the neutral mutation theory of evolution. We can, for example, assume that all sites in the gene change at the same rate μ , according to one of the standard Markov models for base substitution, which specify probabilities of change among the four states A, C, G, and T. If we were to know the genealogy of the copies in detail, statistical estimation of the rates of mutation would be possible, as well as testing of hypotheses about the mutational process. The genealogy is itself the result of statistical process, dependent on N_e , the effective population size. This would be the population size if the population reproduced according to an idealized Wright-Fisher model; as it is, it corrects for some departures from such a model. We could imagine using the genealogy to estimate N_e and test hypotheses about it.

However, we don't know the genealogy. We must therefore integrate over our uncertainty about it. This turns out to confound N_e and μ , and create a large computational problem. In this paper, we will outline the problem, our own Markov Chain Monte Carlo approach, and relate it to the work of Griffiths and Tavaré, who have suggested another Monte Carlo sampling approach. We will also sketch how population growth, migration, recombination, and fossil DNA sequences can be accommodated in our scheme.

2. The Coalescent. It has been known since the work of Sewall Wright, in the 1930's, that if we choose two copies of a gene from a random-mating population, the date of their common ancestor is geometrically distributed, with expectation $2N$ generations. As $2N$ is typically reasonably large, it is also well-approximated by an exponential distribution with that expectation. In 1982 Kingman [1, 2, 3] generalized this to n copies by defining the coalescent process, and proving that the distribution of the genealogies of the n copies converges to it when scaled properly. While Kingman's methods were sophisticated, the resulting distribution is easy to describe and use. This is fortunate, for Kingman's result is fundamental to the analysis of population samples of DNA sequences.

In the coalescent in a population whose size is N , one can sample from the distribution of the possible genealogies of n copies by the following procedure:

1. Set $k = n$ and $T = 0$.
2. Draw a random quantity u_k from an exponential distribution with expectation $4N/(k(k - 1))$.
3. Pick two of the k copies of the gene at random, without replacement.

4. Create a node of the genealogical tree which is the immediate common ancestor of these two copies, and which existed $T + u_k$ generations before the present.
5. Replace these two copies by this common ancestor and set $k = k - 1$.
6. Set $T = T + u_k$.
7. If $k = 1$ we are done. Otherwise return to step 2.

Thus we go back through a series of exponential time intervals, combining randomly chosen pairs of lineages, until we get a complete tree. The expected time to reach the common ancestor of all copies is $4N(1 - 1/n)$ generations. The lineages combine rapidly at first, then more slowly as we go back, and the last two are expected to take $2N$ generations to find their common ancestor, more than half of the expected time. An interesting implication of the coalescent is that a sample of modest size has an excellent chance that its common ancestor will also be the common ancestor of all copies of the gene in the population.

Kingman's coalescent is an approximation, valid when $n^2 \ll N$, but it is in practice extraordinarily accurate. Given the departures that real populations show from any of these idealized models, inaccuracy of Kingman's approximation is the least of our worries. Kingman's coalescent defines the prior distribution of genealogies, and has given its name to the whole area: researchers studying ancestry of samples of genes from populations are said to be working on coalescents.

There are many possible departures from the idealized Wright-Fisher model that underlies Kingman's result, but the coalescent is in effect a diffusion approximation. Many different models of reproduction of single populations will have the same diffusion approximation, and hence the same coalescent process, provided we replace the actual population size N by the appropriate effective population size N_e .

3. Likelihoods. The coalescent gives us a prior distribution of the genealogy G' , which has its intervals expressed in generations. As a product of exponential densities, it is easily written down and easily computed. Its density function is

$$(3.1) \quad f(G'|N_e) = \prod_{k=1}^{n-1} \frac{2}{4N_e} \exp\left(-\frac{k(k-1)}{4N_e} u_k\right)$$

where u_k is the length of the interval during which the genealogy G' has k lineages. If we were able to observe the coalescence intervals u_k , we could estimate N_e . Note that the event that actually occurs brings in a factor of $2/(4N_e)$ rather than $k(k-1)/(4N_e)$ as we know which two lineages have coalesced.

Of course, we do not actually observe coalescence intervals. For most kinds of contemporary data, we can observe only the differences between the members of our sample. For example, for DNA sequences, we can see the number of positions

(sites) at which the molecules differ. That gives us a picture of the coalescence times, but only a clouded picture. We need to make inferences about parameters such as N_e by using a model of the change in the DNA. The notion of a molecular clock provides such a model. We assume a Markov process operating independently at each site in the DNA, with a mutation rate μ . By equating long-term change to mutation, we are implicitly basing ourselves on the neutral mutation model of evolution made famous by Motoo Kimura [4, 5]. We can use a stochastic model of DNA change, and make assumptions of independence of change in different sites and in different lineages, to compute the probability of the observed sequences D given a genealogy G' . I have outlined how to do this [6] and Ziheng Yang, Gary Churchill, and I have more recently shown how to incorporate autocorrelated variation of evolutionary rates from site to site using a Hidden Markov Model approach [7, 8, 9, 10].

We cannot be certain of the genealogy G' . In fact, it is the role of the data to illuminate it, however dimly. To compute the likelihood of the coalescent parameter N_e and the mutation rate μ given the data D , we must integrate over all possible genealogies [11, 12]

$$(3.2) \quad \text{Prob}(D|N_e, \mu) = \int_{G'} f(G'|N_e) \text{Prob}(D|G', \mu).$$

We describe the integration below. The probability of D given G' and μ which appears on the right is the probability calculated by our Markov process model of evolution, the same quantity that is computed in maximum likelihood inference of phylogenies. The quantity μ is a rate of mutation per generation; in more complex cases this may be replaced by several parameters.

Neither of the terms inside the integral in equation (3.2) is hard to compute. The quantity f is given by (3.1) and the other probability requires effort proportional to the total number of DNA bases in our sample, times the square of the number of states at a site, which is 4. The computational problem comes from the vast size of the space of genealogies G' . The space of values of G' is a union of a very large number of Euclidean pieces. Edwards [13] enumerated these: they are his “labelled histories”. With n sequences there are $n!(n-1)!/2^{n-1}$ of them, so that with only 10 sequences there are 2.571×10^9 labelled histories. Each one of these has $n - 1$ node times. The integration in (3.1) must be over all values of these, so that each of these billions of terms integrates over $n - 1$ dimensions. Clearly there is a computational problem here.

All attempts to find mathematical simplifications for this integration have so far failed. Nevertheless two groups – Griffiths and Tavaré and ourselves – have attempted to use Monte Carlo integration. This can work because many of the billions of possible labelled histories make rather little contribution to the integral, because they lead to very low values of the term $\text{Prob}(D|G')$. We will describe our approach first, and then show the relationship between the two approaches, which appear at first sight to be quite different.

4. A Metropolis-Hastings approach. Our approach has been to use Markov Chain Monte Carlo sampling, in particular the Metropolis-Hastings method [14, 15]. We want to sample from the terms of (3.2) using importance sampling, with our importance function being the function that is being integrated. Our approach for the simplest case – a single population of constant size, with no recombination – is outlined by Kuhner et. al. [16, 17].

In that case, it turns out that we can change the time scale of the genealogies. the entities G' have their node times given in generations. Instead we can rescale them to be in units of $1/\mu$ generations, where μ is the underlying neutral mutation rate of the DNA model that we use. Thus if a node in the genealogical tree is 100,000 generations ago, and the underlying mutation rate μ is 10^{-7} , when rescaled the node is 0.01 mutations ago. These are of course expected mutations per site, not actual mutations. informally, we can write this by saying that the genealogy is now G rather than G' , and

$$(4.3) \quad G = \mu G'.$$

The result of this change of variables is of course to alter the density f as well. The coalescence intervals u_k in (3.1) are replaced by $v_k = \mu u_k$, and a factor of $1/\mu$ comes into each term in the resulting density as well. The result is the density:

$$(4.4) \quad g(G|\Theta) = \prod_{k=1}^{n-1} \frac{k(k-1)}{\Theta} \exp\left(-\frac{k(k-1)}{\Theta} v_k\right)$$

where $\Theta = 4N_e\mu$. This resembles closely the widely-used parameter θ that is frequently estimated in evolutionary genetics, except that it contains the neutral mutation rate per site rather than locus.

The result of this change of scale is that the probability $\text{Prob}(D|G', \mu)$ can be replaced by $\text{Prob}(D|G)$, as the branch lengths of G are already multiplied by the mutation rate. In most DNA models, the elapsed time t in generations must be multiplied by a rate of mutation μ before it can be used. If we are given the product μt we can compute the transition probability directly from it. The result is that (3.2) now becomes:

$$(4.5) \quad \text{Prob}(D|\Theta) = \int_G g(G|\Theta) \text{Prob}(D|G).$$

Our objective becomes computing the likelihood of the parameter Θ .

To approximate the integral we take as our importance function the quantity $g(G|\Theta) \text{Prob}(D|G)$, which immediately raises the issue of what value of Θ to use. Ideally one would want to sample at the maximum likelihood value of Θ , but we cannot know in advance what this will be. Our strategy has been to make a rough estimate of Θ , which we call Θ_0 , and use that for an initial sampling, then find a maximum likelihood value based on that first Markov chain. This is then taken as the provisional value for a second Markov chain, and so on. We have usually run 10 of these chains, then two much longer ones at the end. The

final likelihood curve is computed from the second of these long chains. In our programs the user can customize the number and lengths of the chains.

5. Importance sampling and likelihood curves. One useful property of Metropolis-Hastings sampling is that we can estimate the whole likelihood curve from a single run of a Markov chain, rather than having to compute each point on the likelihood surface from a separate run. Suppose that we sample n genealogies from a Markov chain which has probability proportional to $g(G|\Theta_0) \text{Prob}(D|G)$. Call the sampled genealogies the G_i . The usual importance sampling formula for Monte Carlo integration gives:

$$(5.6) \int_G g(G|\Theta) \text{Prob}(D|G) \simeq \frac{1}{n} \sum_{i=1}^n \frac{g(G|\Theta) \text{Prob}(D|G)}{g(G|\Theta_0) \text{Prob}(D|G)} = \frac{1}{n} \sum_{i=1}^n \frac{g(G|\Theta)}{g(G|\Theta_0)}$$

this allows us to estimate the likelihood for other values of Θ from a run of the Markov chain at Θ_0 . Note that the likelihood curve depends only on the Kingman priors of the sampled G_i at Θ and at Θ_0 . This makes it seem that the data are not involved at all; they actually affect the Markov Chain Monte Carlo sampling process and affect the final likelihood through their effect on which G_i are sampled.

6. The Markov Chain sampling. Our samples of the genealogies G must come from a distribution proportional to $g(G|\Theta_0) \text{Prob}(D|G)$. We achieve this through a sampling based on conditional coalescents. A conditional coalescent may be described as a distribution on G that has its density proportional to the coalescent density $g(G|\Theta_0)$ on some domain of G 's, and has density 0 elsewhere. In our programs the conditional coalescents are created by a process of dissolving part of a tree, and reforming that part by allowing lineages to sample their ancestry randomly according to a conditional coalescent. In the original paper by Kuhner et. al. [16], the region of the tree that was dissolved had a single lineage at its base and three lineages at its top. The three lineages, which were not necessarily contemporaneous, were then re-formed into a tree by allowing them to coalesce, but requiring that all three coalesce into a single lineage by the time the base of the dissolved region was reached. The details of how this was done will be found in that paper.

More recently, we have changed to a different conditional coalescent suggested by Peter Beerli. In this, a lineage is selected, and is disconnected from the genealogy, with the lineage being dissolved back up the tree to the next highest coalescent node. It is then allowed to sample its ancestry downwards (backwards in time) until it re-connects to the tree. Note that sometimes this will mean it reconnects below the previous root of the tree. Figure 2 shows this process in a single population. A branch of the tree is chosen at random. In this case it is the one below tip z (tree A). Tree B shows the tree with that branch removed. In tree C we see the process of simulating the conditional coalescence of that lineage with the remaining ones. During the topmost interval of the tree (the time down




FIG. 2. *A conditional coalescent method of altering a tree. A single lineage is chosen at random to be altered (the lineage below z in A). It is removed from the tree (B) and then its coalescence with the remaining lineages is simulated (C). Tree D shows the result.*

to line j), the instantaneous rate of coalescence of that lineage with each of the three others is $2/\Theta$, for a total of $6/\Theta$. We generate an exponential random variate with mean $\Theta/6$, which is the time until coalescence of that lineage with one of the three others. In this case (line 1 in tree C) the time is too long, and takes the lineage past line j . We then consider the lineage to have remained distinct back as far as line j . Starting at that time, we have two other lineages, for a total instantaneous rate of coalescence of $4/\Theta$. We then draw an exponential variate with mean $\Theta/4$. This time, which defines line k , turns out to be a time above the next coalescence, which is the root of the tree. So we connect our new lineage to the tree at the time of line k , choosing one of the two lineages as the one to which it will be connected. The resulting tree is D.

Note that it is possible for any of the lineages, other than the one that is below the root, to be chosen to be dissolved, and it may reconnect to the tree below the original root. The method requires one exponential variate to be generated for each coalescence interval on the remaining part of the tree. If there are m other lineages in an interval, the instantaneous rate of coalescence with them is $2m/\Theta$.

As an example, Figure 3 shows the likelihood curve generated by a run of on the mitochondrial DNA data set of Ward et. al. [18]. The estimate of Θ is 0.0396. Taking an interval two units of log-likelihood below the maximum suggests that the estimate lies between about 0.03 and 0.055. This curve was generated by two long chains of 12,000 steps each, sampling trees every 20 steps. Further details are given by Kuhner et. al. [16].

The method is computationally feasible on workstations or fast desktop computers. Computational effort seems to rise slowly with the number of sequences, especially since we can re-use many of the likelihood computations from one tree to the next. If only part of a tree has changed we can re-use the likelihoods from the rest of the tree. However there are no easy generalizations about how long the Markov chains must be run.

7. The method of Griffiths and Tavaré. Our Monte Carlo sampling approach was preceded by the pioneering and innovative method of Griffiths and Tavaré [19, 20, 21]. At first sight their method appears to bear no relation to ours, and to have considerable advantages over it. A close examination shows that the two methods are related, and makes clear the advantages and disadvantages of our approach.

Griffiths and Tavaré have as their objective the same likelihood function that we compute. They form a system of recurrence equations expressing this likelihood in terms of likelihoods for data sets that have resulted from one fewer evolutionary event. In principle recursive evaluation of these equations, as in an earlier paper by Griffiths [22], will yield the desired likelihood. However, the recursion expands rapidly, and one must therefore use some approximate method of evaluating it. Griffiths and Tavaré [19, 20, 21] choose sample paths down through the recursion randomly. The great advantages of this method are that the computations are rapid, and each such sample path is independent of all the others.




FIG. 3. The log-likelihood curve for Θ for the data of Ward et. al.




FIG. 4. *A history of mutation, coalescences, and ancestral nucleotide choices which is one that could result in a given set of four sequences. Such histories are, in effect, what Griffiths and Tavaré's method samples. The events are described from a point of view looking backwards in time from the present.*

By contrast, our samples are autocorrelated, leading to serious problems knowing how long to continue the sampling. In each of our samples, the likelihood of a tree must be computed. Even if parts of the computation can be re-used, this is much more effort than is needed for their method.

Each step in their sampling goes back one level in the recursion, and amounts to a decision as to what the next most recent event in the genealogy is. The sequence of choices that Griffiths and Tavaré make correspond to a sequence of events in evolution. Going backwards in time, their events are mutations and coalescences, plus choices of the ancestral nucleotides at each site. Figure 4 shows such a history of events, leading to a set of four DNA sequences. It corresponds to one path through their recursion. Note the difference between such a history (H) and the genealogy (G) that we sample. Our genealogy has branch lengths; theirs does not, at least in the simplest case. They specify the place of occurrence of each mutation, while our likelihoods must sum over all possible placements of mutations on the tree. Nevertheless, we can regard their method as Monte Carlo

integration. We can make an equation analogous to our equation 4.3:

$$(7.7) \quad L = \text{Prob}(D|\Theta) = \sum_H \text{Prob}(D|H) \text{Prob}(H|\Theta),$$

where H is a history of events, corresponding to a sequence of choices in Griffiths and Tavaré's recursion. The histories that they sample have the property that they must always lead to the observed sequences. Thus $\text{Prob}(D|H)$ is, trivially, always 1. The term $\text{Prob}(H|\Theta)$ is simply the product of probabilities of the individual events in H . In the history shown in Figure 4, the most recent event could have been a mutation in any of the 11 sites in any of the four sequences, and each could have come from any of three other nucleotides. The particular event that is shown is a coalescence. There are only two sequences (1 and 3) that are identical, and thus could have coalesced at this point. The rate of coalescence for this pair will be $1/(2N_e)$. The next event is a mutation. If we use, for simplicity, a symmetric Jukes-Cantor model of evolution, the rate of occurrence of a particular mutation from C to A at a particular site will be $\mu/3$, where μ is the total mutation rate per site.

Consider all possible histories H , those that lead to the observed sequences as well as those that do not. As the most recent event, there $4 \times 11 \times 3$ possible mutations, and $4 \times 3/2$ possible coalescences. The fraction of this probability contributed by the most recent coalescence in Figure 4 is then $(1/(2N_e)) / (6/(2N_e) + 44\mu)$, which turns out to be $1/(6 + 22\Theta)$. Continuing in this fashion we can calculate the probability of the particular sequence of events in Figure 4 to be

$$\left(\frac{1}{6 + 22\Theta} \right) \left(\frac{\Theta}{18 + 99\Theta} \right) \left(\frac{\Theta}{18 + 99\Theta} \right) \left(\frac{2}{6 + 33\Theta} \right) \left(\frac{1}{1 + 11\Theta} \right) \left(\frac{1}{4} \right)^{11}.$$

The last term is the probability that the initial DNA sequence is as shown in Figure 4. In effect what Griffiths and Tavaré do is to sum over all such histories, adding up this quantity for all those that lead to the observed data.

Griffiths and Tavaré at each stage are considering all possible most recent events that could have led to the observed sequences. They use importance sampling, by sampling at each stage from among the possible events in proportion to their rate of occurrence. Thus at the first stage in the above calculation, they choose among the one possible coalescence and the 33 possible mutations in proportion to the contributions each would make to the numerator (in that case $1/(2N_e)$ versus $\mu/3$). This needs the usual importance sampling correction. Their sampling is done, as ours is too, at a trial value Θ_0 . Suppose that f is the distribution $\text{Prob}(H|\Theta)$, unconditioned on the data, and h is the density from which we sample instead. The importance sampling correction is

$$(7.8) \quad L(\Theta) = \mathcal{E}_f [\text{Prob}(D|H)] = \mathcal{E}_h \left[\frac{f}{h} \text{Prob}(D|H) \right]$$

and since for h we always have $\text{Prob}(D|H) = 1$, the likelihood is just the expectation over h of f/h .

A history H consists of a series of choices. Suppose that history H_i has at stage j a series of possibilities, with the terms of the Griffiths/Tavaré recursion being the $a_{ijk}(\Theta_0)$. Suppose that one that is actually chosen in history H_i has term $b_{ij}(\Theta_0)$. Then the probability of having taken this choice is

$$(7.9) \quad \frac{b_{ij}(\Theta_0)}{\sum_k a_{ijk}(\Theta_0)}$$

and the probability of the history is the product of this over all j , the number of these depending on the number of events in history H_i . This is the expression for h . The distribution f is similar except that it has Θ in place of Θ_0 , and a wider range of possible events, including those which conflict with the data. The full set of events at stage j in this distribution we call the $c_{ijk}(\Theta)$.

We end up with

$$(7.10) \quad L(\Theta) = \mathcal{E}_h \left[\frac{\left(\frac{\prod_j b_{ij}(\Theta)}{\prod_j (\sum_k c_{ijk}(\Theta))} \right)}{\left(\frac{\prod_j b_{ij}(\Theta_0)}{\prod_j (\sum_k a_{ijk}(\Theta_0))} \right)} \right] = \mathcal{E}_h \left[\prod_j \frac{b_{ij}(\Theta)}{b_{ij}(\Theta_0)} \frac{(\sum_k a_{ijk}(\Theta_0))}{(\sum_k c_{ijk}(\Theta))} \right]$$

Griffiths and Tavaré's method consists of sampling from h to approximate this expectation by averaging the ratio on the right. A careful reading of their papers will show that the above expression is precisely what they compute. Thus their method too can be considered a Monte Carlo integration method with an importance function.

Given the independence of their samples, and the rapidity with which they can compute them, one might expect their method to be unequivocally superior to ours. We are, after all, burdened by more computation and autocorrelated samples. The difficulty with their method is that the distribution h from which they sample does not sample from the histories in proportion to their contribution to the likelihood. There is thus some wasted effort. By contrast our Metropolis-Hastings sampling is supposed to sample from genealogies in proportion to their contribution to the likelihood. We thus have reason to hope that our method might do better in some cases. The problem is most easily seen when considering how Griffiths and Tavaré's method will handle two DNA sequences. If those sequences happen to differ by (say) 2 bases, the mutational events that are sampled will include not only the precise changes needed to make the two sequences identical, but also all other changes in all other sites. Thus a great deal of sampling may be needed to sample from the events that contribute most of the likelihood. Griffiths and Tavaré [21] have worried aloud about this very issue.

8. Population growth. The model of an isolated population of constant size can be extended by allowing the population to grow exponentially. Griffiths and Tavaré [19] have done so, and so have we [23]. Our program FLUCTUATE is currently in distribution. In a population of effective size $N_e(t)$ with k lineages, the rate of coalescence is $k(k - 1)/(4N_e(t))$. If the effective size grows exponentially

at rate r , so that when t is the time back from the present (“dual time”),

$$(8.11) \quad N_e(t) = e^{-rt} N_e(0)$$

Taking this into account in the time to coalescence, that density function is [23]

$$(8.12) \quad f(t) = e^{\left[-\frac{k(k-1)}{4N_e(0)r}(e^{rt}-1)\right]} e^{rt} \frac{2}{4N_e(0)}.$$

This can be used to make a counterpart to equation (3.1) straightforwardly. Griffiths and Tavaré [19] have used this for joint likelihood inference of the current value of Θ and the growth rate. We have more recently produced a Hastings-Metropolis algorithm for a similar model.

Once the mutation rate μ is introduced and the branch lengths of the genealogical trees rescaled in units of expected mutations per site, the parameters of the likelihood turn out to be the current value of $4N_e\mu$, called Θ , and the growth rate per unit branch length, which is $g = r/\mu$. The likelihood surfaces in these parameters usually contain long, narrow ridges. At any given value of g , the estimation of Θ has reasonable power, but there is usually a long, narrow, slightly curving ridge whose top is nearly flat. It runs nearly parallel to the g axis, but curving gradually upwards as higher values of g are reached.

There turns out to be surprisingly little power to estimate g , except in cases where the true value of g is large. Even more surprising is the strong bias in the estimate of g . When data sets are generated from a model that has no population growth, they much more often cause us to estimate a large positive g than a negative g . The behavior is so startling as to make us wonder whether it simply be the result of a program bug.

We can verify that the bias is real by using (8.12), and considering the case of two samples ($k = 2$). Suppose that we had very long, nonrecombining sequences. That would allow us to make a precise estimate of the rescaled time $T = \mu t$ to coalescence. The likelihood function can be written in terms of g and $\Theta = 4N_e(0)\mu$.

$$(8.13) \quad \text{Prob}(T|\Theta, g) = e^{\left[-\frac{2}{\Theta g}(e^{gT}-1)\right]} e^{gT} \frac{2}{\Theta}.$$

In the case of two samples, let us assume that Θ is known, and set in (8.13) to its true value, and that we are estimating g . There is no explicit formula solving for the maximum likelihood estimate \hat{g} in terms of T , but the likelihood can be maximized numerically. Now imagine a population whose true growth rate is zero, and whose value of Θ is known to be 1. The scaled coalescent time T for sample size 2 will be distributed exponentially with mean 0.5.

In Figure 5, the maximum likelihood estimate \hat{g} is shown for quantiles of that distribution. It is striking that 87% of the time the estimate is positive, and very strongly positive for small coalescent times (below 0.08 the curve is too high to fit onto this figure). The other 13% of the time the estimate is negative, though only moderately so. The bias in \hat{g} can be seen: it is the average height of the curve, which is strongly positive. Note that the growth rate scale means that




FIG. 5. Estimates of growth rate for $n = 2$ in a data set with a large number of sites, so that coalescence time can be estimated accurately. For a case where Θ is known and the true growth rate is 0, the estimates for different quantiles of the coalescence time are shown. A large bias toward inferring growth is apparent.

$g = 20$ implies growth of the population by a factor of e^{10} during the expected time for two samples to coalesce. Even at the median of the coalescence times, the bias implies that we infer growth by a factor of $e^{2.166}$ during the average coalescence time. As our Metropolis-Hastings algorithm is not used here, this calculation is an independent check of the reality of the bias.

This bias sounds like a serious problem for Monte Carlo integration methods. It is, but we are convinced that it is an equally serious problem for all other methods. However, although the point estimates are biased, if we make interval estimates using the usual chi-square approximation to the distribution of the likelihood ratio, accepting all values of g whose log-likelihood is within 3 units of the peak (in the case where two parameters are being estimated), the true value of 0 is within the interval almost 95% of the time. In addition, the bias becomes much smaller as we add data from more loci [23].

9. Migration. We can also extend the model to allow for multiple populations exchanging migrants. This has been done by Nath and Griffiths [24], who estimate the migration rates for populations whose values of Θ are known. We have [25] extended our Metropolis-Hastings method to a two-population case, to estimate the two values of Θ and two migration rates. This seems to have advantages over methods using statistics like F_{ST} , as those cannot estimate all four parameters independently. An extension to n populations is in progress.

10. Sequential Sampling. In studies of ancient DNA, we have samples that are not contemporaneous. In studies of the course of viral infection in a patient (as in HIV) one may also have sequential samples. The coalescent likelihood approach is readily adapted to such cases [26]. Suppose that we have a genealogical tree G^* whose branch lengths are actual times, with some tips not contemporaneous. Let the generation time of the organism be τ . As one proceeds down the tree, there are two possible events, the entry of a new sample (which has probability 1 at certain known times) or a coalescence. In place of equation 1.1, we have a product of terms, the j -th of which is either

$$(10.14) \quad \frac{2}{4N_e} \exp \left[-\frac{k_j(k_j - 1)}{4N_e} \frac{u_j}{\tau} \right]$$

or

$$(10.15) \quad \exp \left[-\frac{k_j(k_j - 1)}{4N_e} \frac{u_j}{\tau} \right]$$

depending on whether there is a coalescence or a new sample at the bottom end of interval j . Note that k_j is the number of lineages that exist in the genealogy during interval j . Note also that the chronological lengths of the interval have been divided by the generation time to convert them into generation times. The probability of the data given G^* also needs a conversion: it depends on the product of the per-generation mutation rate per site, μ and the generation time elapsed, which is t/τ .

The result is that we can restate equation (3.2) as

$$(10.16) \quad \text{Prob}(D|N_e\tau, \mu/\tau) = \int_{G^*} f(G^*|N_e\tau) \text{Prob}(D|G^*, \mu/\tau).$$

so that the two parameters that can be estimated are $N_e\tau$ and μ/τ . This means that if we know the generation time τ we can estimate N_e and μ separately. Alternatively if (for example) we know μ , we can estimate N_e as well as the generation time τ . Note that the integration over G^* would involve all possible labelled histories and coalescent times, but would not alter the times at which the samples were taken, these being assumed known.

11. Recombination. All of the above cases involve sequences with no recombination. They are thus appropriate for mitochondrial DNA but of dubious value in the nuclear genome. For this reason it has been of great interest to

everyone involved with coalescent likelihood methods to have a way of dealing with recombination. As usual, we have come in second in the race, as Griffiths and Marjoram [27] have an algorithm that infers the likelihood of a sample with two parameters, $4N_e\mu$ and $4N_ec$, where c is the recombination fraction per site. Their method requires substantial computation to adequately sample the histories. Their method makes use of an “ancestral recombination graph” [28] originally described by Hudson [29]. This shows coalescences and recombination events. The latter branch as one goes rootwards, and at each such branching one needs to specify which sites take each of the two routes.

We have also produced a program for inferring these two parameters, [manuscript in preparation]. Although the Metropolis-Hastings approach helps concentrate the sampling on the relevant genealogies, the number of these is so large that the computation is still slow. Figure 6 shows contours of a likelihood surface produced in one of our runs. There are serious problems ahead, as we need to know how long to run the Markov chains to get an accurate answer, and this is generally unknown. However there are also opportunities. One involves using these methods to place a firm likelihood foundation under the widely used genetic mapping method known as linkage disequilibrium mapping.

12. Natural Selection. Until recently it was assumed by everyone that one could not specify the coalescent for sequences that were under natural selection. Only some special cases could be solved, for cases of extreme selection [30, 31, 32]. Recently Neuhauser and Krone [33, 34] made major inroads into the problem, in what are perhaps the best papers on the coalescent since Kingman. They defined a diagram that branches both downward and upward. Unlike the similar diagrams that are produced in cases of recombination, these do not have different alleles following different loops. Instead information flowing upward on the genealogical graph can only pass through certain branches if the genotype contains one of the selected alleles. In the case of recombination, at each site the graph is a tree, although not the same tree at all sites. In the Neuhauser-Krone “ancestral selection graph” the loops are rather more serious. If one tries to compute $\text{Prob}(D|G)$ on them, likelihood must be propagated simultaneously, not independently, around both sides of a loop. However, they were able to define a recursion system that could be evaluated by Griffiths and Tavaré’s method.

Neuhauser and Krone’s work is an enormous and stimulating advance. But it seems ill adapted to our Metropolis-Hastings approach, since for moderate values of $4N_e s$ the number of loops in the ancestral selection graph can become large. Stimulated by it, J.F. has started work on a different method, which involves carrying out Metropolis-Hastings simulation of the frequencies of the selected alleles, as well as the coalescent of other alleles within those alleles, and the “migration” between them that is caused by mutation and recombination. There are no results to report yet.

13. Software Distribution Our package LAMARC (which stands for Likelihood Analysis by Metropolis Algorithm for Random Coalescents) is available




FIG. 6. Contours of the likelihood surface from a single run of a Metropolis-Hastings sampler with recombination on simulated data. The axes are $\Theta = 4N_e\mu$ and the recombination parameter $\rho = 4N_e c$. The contours shown are 1, 2, 3, ... units of log-likelihood below the peak. The true values of the parameters are shown by the dashed lines.

free from its Web site:

<http://evolution.genetics.washington.edu/lamarc.html>

as C source code plus PowerMac and Windows executables. It is readily compiled on workstation C compilers (except for the cc compiler on SunOS systems). As of this writing three programs were in distribution: COALESCE, which analyzes a single population of constant size, FLUCTUATE, which analyzes exponentially growing single populations, and MIGRATE, which analyzes two populations exchanging migrants. More programs and more features will probably be available by the time you read this.

14. An Object-Oriented Fantasy. Even if we could solve some of the problems of how long to run the Markov Chains, the sampling approach has one other serious problem. We like to call it “the 2^8 programs problem”. Each one of these Markov Chain Monte Carlo programs is enormously difficult to write. It takes each of us about 2 years to write and debug one of them. And yet, the present programs are highly limited. We have programs that add one complication (population growth, migration, recombination) but do not combine these in the same program. And yet there are more complications (such as natural selection, speciation, and gene conversion) that need to be considered. Any user may want to pick some particular combination of, say, 8 complications. Do we need to resign ourselves to spending the next 250 years writing all possible programs?

There is one way out. Object-oriented programming methods (such as are embodied in C++ or Objective C) allow a program to self-assemble in response to a user’s requirements. We therefore intend to try to create such an environment. The user would select which combination of evolutionary forces, historical events, genetic situations, and population structure were needed. The program would then use only those classes and subclasses needed to run that particular combination. Thus more like 8 programs than 2^8 need to be written. The issue of chain length remains, and we as yet have no experience with the serious issue of user interface – how do we represent the results of runs that have many parameters, for example?

Nevertheless one may fantasize about an “evolutionary genetics black box”. The user puts in the data and the model, and out come likelihood inferences about the parameters. One still needs to know population genetic theory, of course, to comprehend the model. But a large fraction of the kind of work that has filled theoretical journals in population genetics may become obsolete if this fantasy can be realized. Many papers start with a theoretical model, pose the question of what the expected value of some statistic (such as the probability of monomorphism, or of fixed differences between populations, or the variance of heterozygosity), and after much blood, sweat, and tears arrive at a power series, which usually remains unused by those with data. We may hope that this era can be succeeded by one where the same effort can be redirected to formulating the model and improving the computational methods.

Acknowledgments. We wish to thank Dick Hudson, Robert Griffiths, Simon Tavaré, and Kermit Ritland for helpful communications.

REFERENCES

- [1] Kingman, J. F. C. 1982. The coalescent. *Stochastic Processes and Their Applications* 13: 235-248.
- [2] Kingman, J. F. C. 1982. On the genealogy of large populations. *Journal of Applied Probability* 19A: 27-43.
- [3] Kingman, J. F. C. 1982. Exchangeability and the evolution of large populations. pp. 97-112 in Koch, G. and F. Spizzichino, eds. *Exchangeability in Probability and Statistics. Proceedings of the International Conference on Exchangeability in Probability and Statistics, Rome, 6th-9th April, 1981, in honour of Professor Bruno de Finetti*. North-Holland Elsevier, Amsterdam.
- [4] Kimura, M. 1968. Evolutionary rate at the molecular level. *Nature* 217: 624-626.
- [5] Kimura, M. 1983. *The Neutral Theory of Molecular Evolution*. Cambridge: Cambridge University Press.
- [6] Felsenstein, J. 1981. Evolutionary trees from DNA sequences: a maximum likelihood approach. *Journal of Molecular Evolution* 17: 368-376.
- [7] Yang, Z. 1993. Maximum-likelihood estimation of phylogeny from DNA sequences when substitution rates differ over sites. *Molecular Biology and Evolution* 10: 1396-1401.
- [8] Yang, Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods. *Journal of Molecular Evolution* 39: 306-314.
- [9] Yang, Z. 1995. A space-time process model for the evolution of DNA sequences. *Genetics* 139: 993-1005.
- [10] Felsenstein, J. and G. A. Churchill. 1996. A Hidden Markov Model approach to variation among sites in rate of evolution *Molecular Biology and Evolution* 13: 93-104.
- [11] Felsenstein, J. 1988. Phylogenies from molecular sequences: inference and reliability. *Annual Review of Genetics* 22: 521-565.
- [12] Felsenstein, J. 1992. Estimating effective population size from samples of sequences: inefficiency of pairwise and segregating sites as compared to phylogenetic estimates. *Genetical Research* 59: 139-147.
- [13] Edwards, A. W. F. 1970. Estimation of the branch points of a branching diffusion process. *Journal of the Royal Statistical Society B* 32: 155-174.
- [14] Metropolis, N., A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller. 1953. Equation of state calculations by fast computing machines. *Journal of Chemical Physics* 21: 1087-1092.
- [15] Hastings, W. K. 1970. Monte Carlo sampling methods using Markov chains and their applications. *Biometrika* 57: 97-109.
- [16] Kuhner, M. K., J. Yamato, and J. Felsenstein. 1995. Estimating effective population size and mutation rate from sequence data using Metropolis-Hastings sampling. *Genetics* 140: 1421-1430.
- [17] Kuhner, M. K., J. Yamato, and J. Felsenstein 1997. Applications of Metropolis-Hastings genealogy sampling. pp. 183-192 in *Progress in Population Genetics and Human Evolution*, ed. P. Donnelly and S. Tavaré. IMA Volumes in Mathematics and its Applications, volume 87. Springer Verlag, Berlin.
- [18] Ward, R. H., B. L. Frazier, K. Dew-Jager, and S. Paabo. 1991. Extensive mitochondrial diversity within a single Amerindian tribe. *Proceedings of the National Academy of Sciences, USA* 88: 8720-8724.
- [19] Griffiths, R. C. and S. Tavaré. 1994. Sampling theory for neutral alleles in a varying environment. *Philosophical Transactions of the Royal Society of London, Series B (Biological Sciences)* 344: 403-10.

- [20] Griffiths, R. C. and S. Tavaré. 1994. Ancestral inference in population genetics. *Statistical Science* 9: 307-319.
- [21] Griffiths, R. C. and S. Tavaré 1994. Sampling probability distributions in the coalescent. *Theoretical Population Biology* 46: 131-159.
- [22] Griffiths, R. C. 1989. Genealogical tree probabilities in the infinitely-many-site model. *Journal of Mathematical Biology* 27: 667-680.
- [23] Kuhner, M. K., J. Yamato, and J. Felsenstein. 1998. Maximum likelihood estimation of population growth rates based on the coalescent. *Genetics*, in press.
- [24] Nath, H. B. and R. C. Griffiths. 1996. Estimation in an island model using simulation. *Theoretical Population Biology* 50: 227-253.
- [25] Beerli, P. and J. Felsenstein. 1998. Maximum likelihood estimation of migration rates and effective population numbers in two populations using a coalescent approach. Submitted for publication.
- [26] Rodrigo, A. G. and J. Felsenstein. 1998. Coalescent approaches to HIV-1 population genetics. in press in *Molecular Evolution of HIV*, ed. K. A. Crandall. Johns Hopkins University Press, Baltimore.
- [27] Griffiths, R. C. and P. Marjoram. 1996. Ancestral inference from samples of DNA sequences with recombination. *Journal of Computational Biology* 3: 479-502.
- [28] Griffiths, R. C. and P. Marjoram. 1997. Computational methods for the coalescent. pp. 257-270 in *Progress in Population Genetics and Human Evolution*, ed. P. Donnelly and S. Tavaré. IMA Volumes on Mathematics and Its Applications, volume 87. Springer, New York.
- [29] Hudson, R. R. 1990. Gene genealogies and the coalescent process. *Oxford Surveys in Evolutionary Biology* 7: 1-44.
- [30] Kaplan, N. L., T. Darden, and R. R. Hudson. 1988. The coalescent process in models with selection. *Genetics* 120: 819-829.
- [31] Hudson, R. R. and N. L. Kaplan. 1988. The coalescent process in models with selection and recombination. *Genetics* 120: 831-840.
- [32] Takahata, N. 1990. A simple genealogical structure of strongly balanced allelic lines and trans-species evolution of polymorphism. *Proceedings of the National Academy of Sciences, USA* 87: 2419-2423.
- [33] Neuhauser, C., and S. M. Krone. 1997. The genealogy of samples in models with selection. *Genetics* 145: 519-534.
- [34] Krone, S. M. and C. Neuhauser. 1997. Ancestral processes with selection. *Theoretical Population Biology* 51: 210-237.

DEPARTMENT OF GENETICS
 UNIVERSITY OF WASHINGTON
 BOX 357360
 SEATTLE, WASHINGTON 98195-7360
 JOE@GENETICS.WASHINGTON.EDU